

University of Dundee

On the TRAIL to truth, or on a road to nowhere?

van Steensel, Maurice A. M.

Published in:
Experimental Dermatology

DOI:
[10.1111/exd.13231](https://doi.org/10.1111/exd.13231)

Publication date:
2017

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):
van Steensel, M. A. M. (2017). On the TRAIL to truth, or on a road to nowhere? *Experimental Dermatology*, 26(3), 225-226. <https://doi.org/10.1111/exd.13231>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Received Date : 14-Sep-2016
Accepted Date : 28-Sep-2016
Article type : Commentary from the Editorial Board

On the TRAIL to truth, or on a road to nowhere?

Maurice A.M. van Steensel, MD PhD

Clinical Professor of Genetic Dermatology

School of Medicine and School of Life Sciences

University of Dundee

Senior Principal Investigator

Institute of Medical Biology

Singapore

Honorary Visiting Professor

Lee Kong Chian School of Medicine

Nanyang Technological University

Singapore

Email: maurice.vansteensel@imb.a-star.edu.sg

Abstract

In this issue of Experimental Dermatology, dr. Melnik presents the hypothesis that acne is caused by inappropriate survival of cells in the sebaceous duct and gland. He proposes in particular that TRAIL mediated apoptosis is inhibited by ductal hypoxia. If it works out, this intriguing idea would suggest interesting new therapies.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/exd.13231

This article is protected by copyright. All rights reserved.

In my comment, I offer a critical appraisal – the TRAIL to acne pathogenesis might be rather different from what dr. Melnik would have us believe.

Acne is by far the most common skin disease. In some countries, its incidence amongst adolescents is approaching 100%. The disorder can be very distressing and causes considerable psychological and social morbidity.

Mild disease will spontaneously resolve. Moderate to severe acne, however, requires medical intervention. Several treatments are available. For most, there is no evidence for efficacy or superiority over other options. Examples include photodynamic therapy, pulsed dye laser and azelaic acid. The treatments with proven efficacy have important disadvantages (1). They all take a long time to act and must be used daily. Topical retinoids can induce cumbersome skin irritation. The antibiotics used to treat acne can cause bacteria to become resistant. As an aside, they are effective even in patients with resistant *Propionibacterium acnes* (2). This observation further challenges the already tenuous notion of *P. acnes* having any pathogenic role in acne. The most effective systemic drug, 13-cis retinoic acid (13RA), has undesirable side effects and is highly teratogenic. There are also persistent reports of suicidal ideation associated with its use. These disadvantages preclude 13RA from being prescribed routinely, even if it can cure acne and prevent scarring.

There is thus a clear need for new drugs that can treat acne quickly and have fewer disadvantages. However, no new drug classes have been developed since 13RA was introduced in 1982. This lack of innovation is due to our incomplete

understanding of the molecular events underpinning acne. What are the mechanisms that drive comedogenesis? What causes inflammation and scarring? Despite decades of research, the underlying cellular pathways remain elusive and no novel drug targets have been identified.

A prolific and original thinker, dr. Bodo Melnik has over the past several years outlined interesting new hypotheses on the pathogenesis of acne (3). Even though these remain to be rigorously tested, they have helped to rekindle interest in the fundamentals of acne. In this issue of Experimental Dermatology, dr. Melnik offers up his latest idea (4). Deftly weaving together his previous thinking and published data obtained in seemingly disparate systems, he arrives at the conclusion that inappropriate pro-survival signaling could be an important factor in acne. Specifically, dr. Melnik proposes that hypoxia in the sebaceous duct and increased PI3K/Akt signaling in sebocytes conspire to inhibit TRAIL-mediated apoptosis in the sebaceous gland and associated structures. Abnormal apoptosis might affect the differentiation of infundibular keratinocytes, contributing to comedo formation. Enhanced sebocyte survival could contribute to the altered sebum production implicated in acne. Isotretinoin would then work by enhancing apoptosis, which is consistent with previously published observations (5).

This line of reasoning is interesting and eminently testable. Certainly, inappropriate protection from apoptosis would be an actionable mechanism that is relatively easy to assay in high-throughput screens. That said, key assumptions underlying dr. Melnik's hypothesis could be challenged. For instance, it is assumed that cornification and sebum production both are based on apoptosis, but there is no

good evidence that they are (6). It is also surmised that acne is, at least in part, a sebocyte disorder. However, it is, at heart, a disease of the infundibulum (7). That is where the comedo forms. There is no good evidence that abnormal sebocyte function, as reflected in altered sebum production, has a causal role. Of particular note, comedone formation is well known to be associated with sebaceous gland atrophy (7). This phenomenon conflicts with a key prediction of dr Melnik's hypothesis: the sebaceous gland should become larger due to sebocytes being protected from apoptosis. Of course, one might argue that the pro-survival effect is limited to the infundibulum alone. But in that case, one would have to explain where this limitation comes from. It is also difficult to reconcile a lack of oxygen with the increased keratinocyte proliferation observed in comedones (8).

Finally, positing ductal hypoxia as a trigger mechanism gives rise to an interesting conflict with one of dr. Melnik's previously published hypotheses. Hypoxia inhibits mTORC1 signaling. If the pilosebaceous unit is truly hypoxic (and it might be), a role for increased mTORC1 activity in the pathogenesis of acne seems less likely. That doesn't rule out pathological activation, but this is more typically associated with malignancy (9).

With so many, sometimes conflicting hypotheses ready for falsification, these are exciting times for acne researchers. The road ahead may not be clear, but the trail blazed by dr. Melnik at least gives us some guidance. Here we go!

References

- 1 Zouboulis C C, Bettoli V. Management of severe acne. Br J Dermatol 2015: **172** Suppl 1: 27–36.
- 2 Tan A W, Tan H-H. Acne vulgaris: a review of antibiotic therapy. Expert Opin Pharmacother 2005: **6**: 409–418.
- 3 Melnik B C, Zouboulis C C. Potential role of FoxO1 and mTORC1 in the pathogenesis of Western diet-induced acne. Exp Dermatol 2013: **22**: 311–315.
- 4 Melnik B C. The TRAIL to acne pathogenesis: let's focus on death pathways. Exp Dermatol 2016:
- 5 Melnik B C. Pro-inflammatory sebocyte growth and survival signalling in acne vulgaris are reversed by pro-apoptotic isotretinoin signalling. Exp Dermatol 2016: **25**: 676–677.
- 6 Costanzo A, Fausti F, Spallone G *et al.* Programmed cell death in the skin. Int J Dev Biol 2015: **59**: 73–78.
- 7 Saurat J-H. Strategic Targets in Acne: The Comedone Switch in Question. Dermatology (Basel) 2015: **231**: 105–111.
- 8 Cunliffe W J, Holland D B, Jeremy A. Comedone formation: etiology, clinical presentation, and treatment. Clin Dermatol 2004: **22**: 367–374.
- 9 van Steensel M A M, van Geel M, Badeloe S *et al.* Molecular pathways involved in hair follicle tumor formation: all about mammalian target of rapamycin? Exp Dermatol 2009: **18**: 185–191.